ORIGINAL ARTICLE

Meta-analysis on the safety and efficacy of early oral feeding after total laryngectomy

Xuelian Yi^{1*}, Chunhong Hu¹, Yi Peng¹, Zili Wen², Xue Li¹, Linlin Ye¹, Qingsong Huang^{1*}

¹Department of Otolaryngology Head and Neck Surgery, Chengdu Second People's Hospital, Chengdu, China. ²Department of Oncology, Chengdu Second People's Hospital, Chengdu, China.

*Xuelian Yi and Qingsong Huang contributed equally to this work.

Summary

Purpose: To evaluate the safety and efficacy of early oral feeding (\leq 3 days) and delayed oral feeding (\geq 7 days) following total laryngectomy.

Methods: Relevant literatures on early and delayed oral feeding following total laryngectomy published before January, 2019 were searched in PubMed, EMBASE, Web of Science, Cochrane Library, CNKI and Wanfang Database. Two reviewers were responsible for selecting literatures, extracting data and cross-check. The incidence of pharyngocutaneous fistula (PCF) was evaluated by calculating OR and 95%CI. Difference in length of stay (LOS) of patients undergoing early oral feeding or delayed oral feeding was compared using standardized mean difference (SMD) and 95%CI. Sensitivity analysis and publication bias examination were conducted.

Results: 14 eligible literatures were enrolled, including 1824 patients who underwent total laryngectomy, with 1250 cases

of early oral feeding and 574 cases of delayed oral feeding. The incidence of PCF was similar in patients receiving early oral feeding or delayed oral feeding following total laryngectomy (OR=1.12, 95%CI=0.81-1.54). LOS was shorter in cases of early oral feeding than those of delayed oral feeding (SMD=-0.77, 95%CI=-1.18-0.36). Reliable conclusions were obtained without obvious publication bias.

Conclusions: Early oral feeding following total laryngectomy shortens LOS relative to delayed oral feeding. No significant difference in the incidence of PCF is observed between early oral feeding and delayed oral feeding, suggesting that early oral feeding following total laryngectomy is safe and efficacious.

Key words: total laryngectomy; early oral feeding; pharyngocutaneous fistula; length of stay; meta-analysis

Introduction

Malignant laryngeal tumors are a general term for head and neck malignant tumors originating from the upper and lower glottis and vocal cords. Most of malignant laryngeal tumors are squamous cell carcinoma [1]. Laryngeal cancer mostly affects males, and the high-risk age is about 60 years. In recent years, the onset of laryngeal cancer becomes younger [2]. The major treatments for laryngeal cancer are surgery, adjuvant chemotherapy and radiotherapy [3,4]. With the improvement of economy and living standard, surgery for laryngeal tumors PCF leads to prolonged LOS and delayed oral feed-

are required to preserve the basic functions of the larynx (swallowing, vocalization and breathing) as much as possible [5]. However, laryngeal cancer patients in T4 stage, most of T3 cases, recurrent cases and elderly patients with bad situation need to undergo total laryngectomy [6].

PCF is a common complication after total laryngectomy, with the incidence of 3% to 65% in the past [7]. In recent years, the incidence of PCF is reported to be 9% to 25% [8-10]. The occurrence of

Corresponding author: Qingsong Huang, BM. Department of Otolaryngology Head and Neck Surgery, Chengdu Second People's Hospital, 10 Qingyunan Street, Jinjiang District, Chengdu 610015, China. Tel: +86028-67830994; Email: 1443072662@qq.com

Received: 12/06/2021; Accepted: 01/08/2021



ing time [11]. Previous researches believed that the incidence of PCF reduces with the postponement of oral feeding. Patients undergoing total larvngectomy are intubated with the nasogastric tube for postoperative feeding through nasal feeding tube. Oral feeding is permitted at 8-10 days after surgery. Long-term intubation and nasal feeding pose a great discomfort and cost a huge economic burden [12]. Current studies have demonstrated that oral fluid diet is safe within 48-72 hours after total laryngectomy without elevating the risk of PCF relative to those with oral feeding at 10 days after surgery [13,14]. Aprigliano et al. [15] identified that the placement of nasogastric tube after total laryngectomy increases the risk of PCF. Early oral feeding should be encouraged following total laryngectomy.

Conclusions on the early oral feeding (≤ 3 days) following total laryngectomy and risk of PCF are controversial. This study analyzed relevant literatures on oral feeding following total laryngectomy by extracting their data. We systematically evaluated the safety and efficacy of early oral feeding (≤ 3 days) and delayed oral feeding (≥ 7 days). Our conclusions provide reliable and convincing evidence of early oral feeding in clinical application.

Methods

Searching strategy

Relevant literatures on early and delayed oral feeding following total laryngectomy published before January, 2019 were searched in PubMed, EMBASE, Web of layed oral feeding.

Science, Cochrane Library, CNKI and Wanfang Database. Key words were as follows: "oral feeding" or "oral intake" and "laryngectomy" or "laryngeal neoplasm". Inclusive criteria were applied: 1) Surgical treatment must be the total laryngectomy; 2) Tumors originating in pharynx and larynx; 3) Study design, experimental methods, statistical analyses and monitoring indexes were similar; and 4) the number of PCF cases and LOS of patients with early oral feeding or delayed oral feeding were provided. Exclusive criteria were applied: 1) Tumors originating in other places except for pharynx and larynx; 2) Surgical treatments were partial laryngectomy and vertical hemilaryngectomy; and 3) repeated studies, low-quality studies and those with incomplete data.

Data extraction

Data acquisition was independently carried out by two reviewers, and a third reviewer was responsible for re-evaluating disagreements. Baseline data acquisition included: first author, study type, sample size, year of publication, monitoring indexes, etc. Procedures for screening literatures were as follows: 1) Exclude ineligible literatures by reviewing titles and abstracts; 2) Exclude ineligible literatures by reviewing the full-text; and 3) for uncertain literatures, supplementary information were reviewed or contacting the author for in-depth information.

Outcome evaluation

(1) Safety (incidence of PCF): PCF is a common and serious complication following total laryngectomy. The safety of early and delayed oral feeding was evaluated by analyzing the incidence of PCF (the ratio of PCF cases and total cases). (2) Efficacy (LOS): The average LOS was calculated for evaluating the efficacy of early and delayed oral feeding.

Table 1. Main characteristics of studies included in the meta-analysis

Year	Study	Nationality	Study design	Numbers of patients		Study index	Study quality
				Early oral intake	Late oral intake		
2018	Huang	China	RD	18	34	PCF	7 ª
2016	Suslu	Turkey	RD	582	20	PCF	6ª
2015	Serbanescu-Kele	Netherlands	RD	36	30	PCF, LOS	7ª
2014	Sousa	Brazil	RCT	44	45	PCF	5 ^b
2014	Timmermans	Netherlands	RD	107	140	PCF	6 ^a
2009	Aswani	South Africa	RD	40	39	PCF, LOS	5ª
2006	Prasad	India	RD	40	38	PCF, LOS	6 ^a
2003	Song	China	RD	21	21	PCF, LOS	6ª
2003	Seven	Turkey	RCT	32	33	PCF, LOS	4^{b}
2001	Volling	Germany	RCT	21	21	PCF	4^{b}
2001	Medina	American	RD	20	18	PCF, LOS	7ª
1998	Soylu	Turkey	RD	252	43	PCF	8ª
1995	Rodriguez-Cuevas	Mexico	RCT	18	17	PCF	5^{b}
1989	Boyce	American	RD	19	75	PCF	7 ª

RD: Retrospective Design; RCT: Randomized Controlled Trials; PCF: Pharyngocutaneous fistula; LOS: Length of stay; a: NOS Scale; b: Jadad Scale

Quality assessment

Randomized controlled trials (RCT) were analyzed using the modified Jadad scale for literature quality evaluation, with 1-3 scores for low quality and 4-7 for high quality. Retrospective studies were analyzed using the Newcastle-Ottawa Scale (NOS) for quality evaluation (>5 score considered as high quality). Only high-quality literatures were enrolled in this study.

Statistical analysis

Statistical analysis was performed using Stata 14 software. The incidence of PCF was evaluated by calculating OR and 95%CI. Difference in LOS of patients undergoing early oral feeding or delayed oral feeding was compared using standardized mean difference (SMD) and 95%CI. The heterogeneity among literatures was tested by Cochran Q test. p<0.05 considered statistically

significant. Fixed-effect model was used when $I^2 < 50\%$ and p>0.05; Otherwise, the random-effects model was used. Begg's funnel plots were depicted for evaluating the publication bias.

Results

Studies characteristics

There were 353 relevant literatures searched and 27 were enrolled after reading the titles and abstracts. Based on the inclusive criteria, finally 14 literatures involving 1250 cases of early oral feeding and 574 cases of delayed oral feeding were enrolled [14,16-28]. Baseline characteristics of these literatures were displayed in Table 1. Report searching and selection process were depicted in Figure 1.





Table 2	2.	Meta-	analys	sis	for	two	gro	ups
---------	----	-------	--------	-----	-----	-----	-----	-----

Study Index	Trials	Heterogeneity		Model	Pooled statistics		
		р	I ² (%)	_			
PCF	14	0.772	0	Fixed	OR (95% CI)=1.12 (0.81,1.54)		
LOS	6	0.003	72	Random	SMD (95% CI) =-0.77 (-1.18,-0.36)		

PCF: Pharyngocutaneous fistula; LOS: Length of stay



Figure 2. Forest plots of merged analyses of pharyngocutaneous fistula.



Figure 3. Forest plots of merged analyses of length of stay.

Safety of early oral feeding and delayed oral feeding following total laryngectomy

The incidence of PCF was recorded for evaluating the safety of early or delayed oral feeding. No significant difference in the heterogeneity was identified among these studies (p=0.772, $I^2=0\%$) (Table 2). Therefore, the fixed-effect model was utilized. The results showed that the pooled OR value was 1.12 (95%CI=0.81-1.54), suggesting that early oral feeding did not increase the risk of PCF relative to delayed oral feeding (Figure 2).

Efficacy of early oral feeding and delayed oral feeding following total laryngectomy

LOS was recorded for evaluating the efficacy of early or delayed oral feeding. There were 6 RCTs recoding the LOS, with 189 cases of early oral feeding and 179 cases of delayed oral feeding. A remarkably heterogeneity was identified among these studies (p=0.003, I²=72%). Therefore,

the fixed-effect model was utilized. The results showed that the pooled SND was -0.77 (95%CI=-1.18-0.36), suggesting a shorter LOS in cases of early oral feeding relative to those of delayed oral feeding (Figure 3).

Sensitivity analysis

Sensitivity analysis was performed for illustrating the impact of individual literature on the summary estimate. As Figure 4 illustrated, sensitivity analysis indicated the final conclusion was not influenced by an individual literature. We believed that our meta-analysis results were robust and stable.

Publication bias

Publication bias in this study was assessed by depicting Begg's funnel plots. The systematic shape of funnel diagram indicated no significant publication bias (Figure 5).



Figure 4. (A) Effect of individual studies on the pooled OR for pharyngocutaneous fistula; **(B)** Effect of individual studies on the pooled SMD for length of stay.



Figure 5. Begg's funnel plot of publication bias test for pharyngocutaneous fistula (A) and length of stay (B).

Discussion

To avoid PCF after laryngeal cancer surgery, postoperative nasogastric tube diet is generally required for about 10 days. The most vulnerable period of the wound is about a week after surgery. Hence, Applebaum et al. [29] proposed that oral feeding should delay to the 10th day after surgery. Recently, clinical observations considered that early oral feeding at day 2-3 after surgery dose not influence the healing and the incidence of PCF [14,30]. This study summarized and analyzed 14 high-quality literatures involving 1824 cases undergoing total laryngectomy. Our conclusion may provide clinical evidences for the application of early oral feeding.

This study demonstrated no significant difference in the incidence of PCF between cases of early or delayed oral feeding. Hence, early oral feeding was proved to be safe, which decreased psychological burden of affected patients. Meanwhile, early oral feeding shortened LOS, which decreased economical burden. We encouraged early oral feeding in the case of no fistula formation, and it could elevate the bed usage rate and reduce medical expense.

Although Apriglano et al. [15] suggested no benefit of early oral feeding following total larvngectomy, they believed that the intubation of nasogastric tube is one of the reasons for PCF. Early oral feeding avoids the long-term usage of nasogastric tube. Medina et al. [21] indicated that early oral feeding does not increase the incidence of PCF, and the avoidance of nasogastric tube usage could relieve the fear and discomfort. Patients undergoing early oral feeding could be more cooperated, leading to a better outcome. Akyol et al. [31] analyzed a total of 110 patients undergoing total laryngectomy in 1986-1992. They considered that no application of nasogastric intubation will not elevate the incidence of PCF, and oral feeding could be applied within the 1-2 days after surgery. Soylu et al. [26] reported that the oral feeding on the third day does not increase the risk of PCF in 295 cases undergoing total laryngectomy. They demonstrated that the third day after total laryngectomy is the most suitable time point for oral feeding. Identi-

cally, Timmemans et al. [23] also encouraged the early oral feeding. This study proposed the safety of early oral feeding as well.

LOS is a crucial marker for evaluating treatment efficacy. Medina et al. [21] showed the shorter LOS in cases of early oral feeding compared with those of delayed oral feeding (7.05 days *vs.* 11.87 days). Song et al. [28] revealed that early oral feeding reduces 4.3 days of LOS in patients undergoing total laryngectomy. Our study also pointed out shorter LOS in cases of early oral feeding. However, Seven et al. [17] illustrated that early oral feeding dose not shorten the LOS.

Sensitivity analysis is conducted to analyze the data using different statistical methods and models, so as to rule out low-quality studies. In this study, sensitivity analysis indicated the reliable and robust conclusion. Publication bias is evaluated according to the symmetry of funnel plots and the concentration to the midline. Here, the systematic shape of funnel diagram indicated no significant publication bias.

Several limitations in this study should be noteworthy: 1) Sample size was relatively small, and further explorations should be conducted with a large sample size to improve the accuracy; 2) Subgroup analyses on the influence of early oral feeding on the onset time of PCF were lacked; 3) Confounding factors of PCF should be taken into consideration, such as radiotherapy, TNM stage of tumors, albumin level and Hb level; and 4) different physicians with varied surgical experiences could influence the conclusion.

Conclusions

Early oral feeding following total laryngectomy shortens LOS relative to delayed oral feeding. No significant difference in the incidence of PCF is observed between early oral feeding and delayed oral feeding, suggesting that early oral feeding following total laryngectomy is safe and efficacious.

Conflict of interests

The authors declare no conflict of interests.

References

- 1. Tan J, Jing YY, Han L et al. MicroRNA-203 inhibits invasion and induces apoptosis of laryngeal cancer cells via targeting LASP1. Eur Rev Med Pharmacol Sci 2018;22:6350-7.
- 2. Steuer CE, El-Deiry M, Parks JR, Higgins KA, Saba

NF. An update on larynx cancer. CA Cancer J Clin 2017;67:31-50.

 Karamanou M, Markatos K, Lymperi M, Agapitos E, Androutsos G. A historical overview of laryngeal carcinoma and the first total laryngectomies. J Buon 2017;22:807-11.

2025

- Dragicevic D, Andelic B, M JR, Kljajic V, Vlaski L, Savovic S. Clinical stage of laryngeal carcinoma and lost time at the moment of diagnosis with 15-year-long interval. Are there any changes? J Buon 2019;24:2041-8.
- 5. Kolator M, Kolator P, Zatonski T. Assessment of quality of life in patients with laryngeal cancer: A review of articles. Adv Clin Exp Med 2018;27:711-5.
- 6. Mannelli G, Lazio MS, Luparello P, Gallo O. Conservative treatment for advanced T3-T4 laryngeal cancer: meta-analysis of key oncological outcomes. Eur Arch Otorhinolaryngol 2018;275:27-38.
- 7. Redaelli DZL, Ferrari L, Tomenzoli D, Premoli G, Parrinello G, Nicolai P. Postlaryngectomy pharyngocutaneous fistula: incidence, predisposing factors, and therapy. Head Neck 1999;21:131-8.
- Busoni M, Deganello A, Gallo O. Pharyngocutaneous fistula following total laryngectomy: analysis of risk factors, prognosis and treatment modalities. Acta Otorhinolaryngol Ital 2015;35:400-5.
- Benson EM, Hirata RM, Thompson CB et al. Pharyngocutaneous fistula after total laryngectomy: a singleinstitution experience, 2001-2012. Am J Otolaryngol 2015;36:24-31.
- 10. Saki N, Nikakhlagh S, Kazemi M. Pharyngocutaneous fistula after laryngectomy: incidence, predisposing factors, and outcome. Arch Iran Med 2008;11:314-7.
- 11. Esteban F, Delgado-Rodriguez M, Mochon A, Solano J, Soldado L, Solanellas J. [Study of in-patient hospital stay following total laryngectomy: multivariable retro-spective analysis of a 442 total laryngectomies]. Acta Otorrinolaringol Esp 2006;57:176-82.
- 12. Gooi Z, Richmon J. Long-term oral intake through a salivary bypass tube with chronic pharyngocutaneous fistula. Am J Otolaryngol 2012;33:762-3.
- 13. Kishikova L, Fleming JC. Oral feeding following laryngectomy: early or delayed? Int J Surg 2014;12:1137-40.
- 14. Suslu N, Sefik HA. Early oral feeding after total laryngectomy: Outcome of 602 patients in one cancer center. Auris Nasus Larynx 2016;43:546-50.
- 15. Aprigliano F. Use of the nasogastric tube after total laryngectomy: is it truly necessary? Ann Otol Rhinol Laryngol 1990;99:513-4.
- Volling P, Singelmann H, Ebeling O. [Incidence of salivary fistulas in relation to timing of oral nutrition after laryngectomy]. Hno 2001;49:276-82.
- 17. Seven H, Calis AB, Turgut S. A randomized controlled

trial of early oral feeding in laryngectomized patients. Laryngoscope 2003;113:1076-9.

- 18. Sousa AA, Porcaro-Salles JM, Soares JM et al. Does early oral feeding increase the likelihood of salivary fistula after total laryngectomy? J Laryngol Otol 2014:1-7.
- 19. Serbanescu-Kele CM, Halmos GB, Wedman J, van der Laan BF, Plaat BE. Early feeding after total laryngectomy results in shorter hospital stay without increased risk of complications: a retrospective case-control study. Clin Otolaryngol 2015;40:587-92.
- 20. Aswani J, Thandar M, Otiti J, Fagan J. Early oral feeding following total laryngectomy. J Laryngol Otol 2009;123:333-8.
- 21. Medina JE, Khafif A. Early oral feeding following total laryngectomy. Laryngoscope 2001;111:368-72.
- 22. Prasad KC, Sreedharan S, Dannana NK, Prasad SC, Chandra S. Early oral feeds in laryngectomized patients. Ann Otol Rhinol Laryngol 2006;115:433-8.
- 23. Timmermans AJ, Lansaat L, Kroon GV, Hamming-Vrieze O, Hilgers FJ, van den Brekel MW. Early oral intake after total laryngectomy does not increase pharyngocutaneous fistulization. Eur Arch Otorhinolaryngol 2014;271:353-8.
- 24. Boyce SE, Meyers AD. Oral feeding after total laryngectomy. Head Neck 1989;11:269-73.
- 25. Rodriguez-Cuevas S, Labastida S, Gutierrez F, Granados F. Oral feeding after total laryngectomy for endolaryngeal cancer. Eur Arch Otorhinolaryngol 1995;252:130-2.
- 26. Soylu L, Kiroglu M, Aydogan B et al. Pharyngocutaneous fistula following laryngectomy. Head Neck 1998;20:22-5.
- 27. Huang N, Zhu YM, An CM et al. [Primary research of early oral feeding after total laryngectomy]. Zhonghua Er Bi Yan Hou Tou Jing Wai Ke Za Zhi 2018;53:428-31.
- 28. Song J, Jing S, Shi H. [The clinical observation of early oral feeding following total laryngectomy]. Lin Chuang Er Bi Yan Hou Ke Za Zhi 2003;17:527-8.
- 29. Applebaum EL, Levine HL. Pharyngeal reconstruction after laryngectomy. Laryngoscope 1977;87:1884-90.
- 30. Kishikova L, Fleming JC. Oral feeding following laryngectomy: early or delayed? Int J Surg 2014;12:1137-40.
- Akyol MU, Ozdem C, Celikkanat S. Early oral feeding after total laryngectomy. Ear Nose Throat J 1995;74:28-30.